

FORM PTO-1390 (REV. 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				0230-0161P	
				U.S. APPLICATION NO. (If known, see 37 CFR 1.5)	
				09/856718	
INTERNATIONAL APPLICATION NO.		INTERNATIONAL FILING DATE		PRIORITY DATE CLAIMED	
PCT/JP99/06616		November 26, 1999		November 27, 1998	
TITLE OF INVENTION					
LAK ACTIVITY ENHANCERS DERIVED FROM EXTRACT OF LENTINUS EDODES MYCELIUM AND LAK *					
APPLICANT(S) FOR DO/EO/US					
ASANO, Kenji; MATSUDA, Yukiko and TAJIMA, Yutaka					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<p>1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1).</p> <p>4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31).</p> <p>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))</p> <p>6. <input checked="" type="checkbox"/> <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau).</p> <p>7. <input checked="" type="checkbox"/> has been transmitted by the International Bureau. WO 00/32212</p> <p>8. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>9. <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).</p> <p>10. <input checked="" type="checkbox"/> <input type="checkbox"/> is transmitted herewith.</p> <p>11. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4)</p> <p>12. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).</p> <p>13. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).</p> <p>14. <input type="checkbox"/> have been transmitted by the International Bureau.</p> <p>15. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p>16. <input checked="" type="checkbox"/> have not been made and will not be made.</p> <p>17. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</p> <p>18. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</p> <p>19. <input checked="" type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</p>					
Items 11. to 20. below concern document(s) or information included:					
<p>21. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98-1.449 and International Search Report w/ cited references</p> <p>22. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</p> <p>23. <input checked="" type="checkbox"/> A FIRST preliminary amendment.</p> <p>24. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.</p> <p>25. <input type="checkbox"/> A substitute specification.</p> <p>26. <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>27. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825.</p> <p>28. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4).</p> <p>29. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).</p> <p>30. <input checked="" type="checkbox"/> Other items or information:</p> <p>PCT/ISA/210</p> <p>PCT/IEPA/409</p> <p>PCT/RO/101</p> <p>Verification of a translation</p>					
*ACTIVITY-ENHANCING FORMULATIONS CONTAINING THE EXTRACT					

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER	
09/856718		PCT/IP99/06616		0230-0161P	
21. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. \$1,000.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. \$710.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	860.00
				\$	130.00
CLAIMS		NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims		13 - 20 =	0	X \$18.00	\$ 0
Independent Claims		1 - 3 =	0	X \$80.00	\$ 0
MULTIPLE DEPENDENT CLAIM(S) (if applicable)		Yes		+ \$270.00	\$ 270.00
TOTAL OF ABOVE CALCULATIONS =				\$	1,260.00
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	0
SUBTOTAL =				\$	1,260.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	0
TOTAL NATIONAL FEE =				\$	1,260.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	0
TOTAL FEES ENCLOSED =				\$	1,260.00
				Amount to be:	
				refunded	\$
				charged	\$

a. ☒ A check in the amount of \$ 1,260.00 to cover the above fees is enclosed.


b. ☐ Please charge my Deposit Account. No. _____ in the amount of \$ _____ to cover the above fees.
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 02-2448.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

Send all correspondence to:
 Birch, Stewart, Kolach & Birch, LLP or Customer No. 2292
 P.O. Box 747
 Falls Church, VA 22040-0747
 (703)205-8000

Date: May 25, 2001

By 
 Gerald M. Murphy, Jr., #28,977

VERIFICATION OF A TRANSLATION

I, the below named translator, hereby declare that:

My name and post office address are as stated below;

That I am knowledgeable in the English language and in the language in which the below identified application was filed, and that I believe the English translation of International Application No. PCT/JP99/06616 is a true and complete translation of the above identified International Application as filed.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Dated this 22nd day of May, 2001

Full name of the translator:

Norihiro FUKAZAWA

Signature of the translator:



Post Office Address:

c/o YUASA AND HARA, Section 206,
New Ohtemachi Bldg., 2-1,
Ohtemachi 2-chome, Chiyoda-ku,
Tokyo, JAPAN

09856718.012802

09/856718

JC18 Rec'd PCT/PTO 25 MAY 2001

PATENT
0230-0161P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: ASANO, Kenji et al. Conf.:
Int'l. Appl. No.: PCT/JP99/06616
Appl. No.: NEW Group:
Filed: May 25, 2001 Examiner:
For: LAK ACTIVITY ENHANCERS DERIVED FROM
EXTRACT OF LENTINUS EDODES MYCELIUM
AND LAK ACTIVITY-ENHANCING
FORMULATIONS CONTAINING THE EXTRACT

PRELIMINARY AMENDMENT

BOX PATENT APPLICATION

Assistant Commissioner for Patents
Washington, DC 20231

May 25, 2001

Sir:

The following Preliminary Amendments and Remarks are respectfully submitted in connection with the above-identified application.

AMENDMENTS

IN THE SPECIFICATION:

Please amend the specification as follows:

Before line 1, insert --This application is the national phase under 35 U.S.C. § 371 of PCT International Application No. PCT/JP99/06616 which has an International filing date of November 26, 1999, which designated the United States of America and was not published in English.

09/856718-01/2001

IN THE CLAIMS:

Please amend the claims as follows:

9. (Amended) The LAK activity-enhancing formulation of claim 4, wherein said formulation is used for treating tumor and/or cancer.

10. (Amended) A method for treating tumor and/or cancer comprising administering an effective amount of the LAK activity-enhancing formulation of claim 4.

REMARKS


The specification has been amended to provide a cross-reference to the previously filed International Application. The claims have also been amended to delete improper multiple dependencies and to place the application into better form.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By

 #3288
Gerald M. Murphy, Jr., #28,977

GMM/rem
0230-0161P

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

(Rev. 02/12/01)

J618 Rev'd PCT/PTO 25 MAY 2001

Docket No.0230-0161 0230-0161P

VERSION WITH MARKINGS TO SHOW CHANGES MADE

The claims have been amended as follows:

9. (Amended) The LAK activity-enhancing formulation of[any one of claims 4 to 8] claim 4, wherein said formulation is used for treating tumor and/or cancer.

10. (Amended) A method for treating tumor and/or cancer comprising administering an effective amount of the LAK activity-enhancing formulation of[any one of claims 4 to 8] claim 4.

SPECIFICATION

LAK ACTIVITY ENHANCERS DERIVED FROM
EXTRACT OF LENTINUS EDODES MYCELIUM AND
LAK ACTIVITY-ENHANCING FORMULATIONS CONTAINING THE EXTRACT

5 FIELD OF THE INVENTION

The present invention relates to the immunological field. In particular, it relates to immune activity enhancers and therapeutic formulations containing an extract of Lentinus edodes mycelium. And more particularly,
10 it relates to materials useful for enhancing LAK activity and LAK activity-enhancing formulations containing the extract.

PRIOR ART

15 The activity of the immune system can be classified into two modes of action: Cell-mediated immunity and humoral immunity. Humoral immunity refers to an immune response involving antibodies, while cell-mediated immunity refers to an immune response produced by cells acting
20 directly on a target.

Lymphocytes play the main role in this cell-mediated immune response and often target cell surface substances as antigens. Cell-mediated immunity is thought to be also involved in immune response against tumor, intracellular
25 parasites, and viruses, transplantation, some drug hypersensitivity and some autoimmune diseases. Generally, the reaction is often specific, but can also be non-specific.

For example, it is known in tumor immunology that tumor cells possess tumor antigens. That is, it is known that there are two types of antigens, one type consisting of tumor-specific antigens (TSA) and the other of tumor-associated antigens (TAA) which is also present in normal cells at a very low level, and upregulated with malignant transformation.

These tumor antigens are expressed by alteration in a genetic sequence or by mutation of autologous cells. A common therapy used against tumor cells having abnormal antigenic expression is immunotherapy, including immunization with a tumor antigen or administration of a drug which enhances the immune function. It is generally known that natural killer (NK) cells demonstrate higher tumor cell-destroying activity than normal cells and that immunotherapy can also enhance the activity of NK cells. NK cells are cytotoxic lymphocytes also present in normal individuals and are known to exhibit MHC antigen-nonrestricted cytotoxicity against tumor cells, virus-infected cells or the like. However, it has been recently shown that tumor cells exist which are resistant to even NK cells.

Dr. S. Rosenberg of the United States National Cancer Institute (NCI) found that incubation of peripheral lymphocytes with interleukin 2 (IL-2) can induce killer cells showing cytotoxicity to a wide range of target cancer cells, including autologous cancer cells, and that these killer cells can kill even NK cell-resistant cancer cells

anemia and thrombocytopenia, as well as causing in vitro damages to lymphocytes, macrophages and vascular endothelial cells. Moreover, IL-2 administered via the oral route is poorly absorbed and must therefore mainly be
5 administered via injection and direct administration at the present time.

Thus, there is a demand for the development of therapeutic agents capable of overcoming these disadvantages, especially agents that could enhance the LAK
10 activity of lymphocytes without using IL-2 or pharmaceutical agents containing such agents.

Anticancer agents are generally targeted at abnormal growth of cancer cells and can be classified by target into nucleic acid biosynthesis inhibitors such as alkylating
15 agents, nucleic acid biosynthesis substrate analogs, antibiotics, steroid hormones, etc. and mitotic inhibitors such as plant alkaloids.

However, such anticancer agents also cause noticeable side effects with regard to normal proliferative cells such
20 as bone marrow, gastrointestinal epithelium, and hair follicle. That is, they generally cause nausea, vomiting, ulcers of the mouth and small bowel, diarrhea, epilation, bone marrow inhibition leading to underproduction of major blood components, and so on, irrespective of the
25 administration route.

Some bacteria, foods and other naturally occurring substances are known to have anticancer properties. Bacteria and food-tye substance are generally far

preferential for use as anti-cancer agents due to their generally benign nature and low side-effect profile. Many attempts have been made to cure cancer by using bacteria, as shown in reports relating to Coley's toxin consisting of
5 a culture filtrate of *Serratia marcescens* and *Streptococcus pyogenes* (1964); treatment of leukemia with BCG (Mathe, G., *Adv. Cancer Res.*, 14, 1, 1971); tumor regression in guinea pigs (Zbar, B., et al., *J. Natl. Cancer Inst.*, 48, 831, 1971); and effectiveness of administration of yeast cell
10 wall polysaccharide against transplanted tumor cells such as sarcoma 180, for example.

Especially, a great amount of research has been conducted into the anticancer effect of polysaccharides derived from yeast such as yeast glucan and yeast mannan,
15 from other bacterium, from lichens and from basidiomycetes. Among them, commercial products currently available on the market as anticancer immunopotentiators include Krestin derived from the cultured mycelia of kawaratake (*Coriolus versicolor*, Basidiomycetes: Polyporaceae) (booster of
20 immune function of hosts, Kureha Chemical Industry and Sankyo Co.Ltd.), a polysaccharide derived from shiitake (*Lentinus edodes*) called lentinan and a polysaccharide derived from suehirotake (*Schizophyllum Commune*).

Lentinus edodes (Shiitake) is a common edible
25 mushroom found in both Japan and China, and has been cultivated in Japan for around 300 years. It has been recently elucidated for its pharmacological effects and effective ingredients and reported to have various effects,

such as growth inhibition effects on transplanted tumor cells in the large bowel and liver in rats and mice (Sugano N. et al., Cancer Letter, 27:1, 1985; Suzuki Y. et al., Journal of the Japan Society of Coloproctology, 43:178, 1990); mitogenic effect (Tabata T. et al., Immunopharmacology, 24:57, 1992; Hibino et al., Immunopharmacology, 28:77, 1994), etc.

DISCLOSURE OF THE INVENTION

10 The present inventors researched the immunopotentiating activity, antitumor activity and/or anticancer activity of Lentinus edodes with a view to providing an inexpensive immunotherapeutic agent having few side effects.

15 The present invention was accomplished on the basis of the finding that an extract of the mycelium, which is a precursor to the edible fruiting body of Lentinus edodes, has a far higher immunopotentiating activity, antitumor activity and/or anticancer activity than the fruiting body
20 and that said extract also has an LAK activity-enhancing activity.

 According to the first aspect of the present invention, therefore, an LAK activity enhancer containing an extract of Lentinus edodes mycelium is provided.

25 "LAK activity" refers to the cytotoxic activity of cytotoxic T-lymphocytes, which attacks tumors unrecognizable by lymphocytes having NK activity but which have little influence on autologous normal cells. "LAK

activity-enhancing" refers to the effect of enhancing this LAK activity, that is directly or indirectly inducing the production of LAK cells from lymphocytes or further enhancing the activity of existing LAK cells.

5 Enhancement of LAK activity increases antitumor activity of LAK cells, which leads to an improvement in the function of the cell-mediated immune system. Thus, the present invention can be applied not only to treatments for improving antitumor activity but also to treatments for
10 improving the immune function.

 Lymphocytes derived from peripheral blood may be treated with LAK activity enhancers of the present invention to enhance LAK activity. They may directly act on lymphocytes derived from peripheral blood preferably at
15 an extract of Lentinus edodes mycelium concentration of 1 μ g or more per 10^6 lymphocytes or 1 μ g or more per 1 ml of peripheral blood. LAK activity enhancers of the present invention may be used in LAK therapy in place of IL-2.

 LAK activity enhancers of the present invention may
20 consist of an extract of Lentinus edodes mycelium on its own or may further contain materials other than an extract of Lentinus edodes mycelium. The range of such materials is not limited so long as the LAK activity enhancing-
25 activity of the extract of Lentinus edodes mycelium is not affected by their use.

 LAK activity enhancers of the present invention may also be combined with other materials having an immunopotentiating activity and antitumor activity and/or

anticancer activity.

According to the second aspect of the present invention, an LAK activity-enhancing formulation comprising an LAK activity enhancer containing an extract of Lentinus edodes mycelium is provided.

LAK activity-enhancing formulations of the present invention may be the extract of Lentinus edodes mycelium itself or pharmaceutical or veterinary formulations comprising an LAK activity enhancer containing an extract of Lentinus edodes mycelium and a pharmaceutically acceptable carrier.

LAK activity-enhancing formulations of the present invention may be suitable for oral administration or injection or percutaneous administration.

LAK activity-enhancing formulations of the present invention may be in the form of a food, drink or feed.

LAK activity-enhancing formulations of the present invention may be used for treating tumor and/or cancer. LAK activity-enhancing formulations of the present invention do not specifically attack specific tumor cells, because they induce LAK cells. Therefore, LAK activity-enhancing formulations of the present invention can be used for treating various tumors, either epithelial or non-epithelial.

LAK activity-enhancing formulations of the present invention may also be used for treating other diseases than tumor or cancer, such as the improvement of the immune function. Specifically, LAK activity-enhancing

formulations of the present invention may also be used for the improvement of immune function compromised by immunocompromising therapies or accidents such as radiation, autoimmune diseases, infectious diseases, etc.

5 LAK activity-enhancing formulations of the present invention can also be used as prophylactic agents against various diseases. For example, they may be administered to prevent tumor and/or cancer or to prevent bacterial or viral infections.

10 LAK activity-enhancing formulations of the present invention may be used in combination with other drugs such as antitumor and/or anticancer agents such as immunopotentiators, antibiotics, analgesics, antiinflammatory agents, etc.

15 According to the third aspect of the present invention, a method for treating tumor and/or cancer comprising administering an effective amount of said LAK activity-enhancing formulation is provided.

20 The method for treating tumor and/or cancer of the present invention may be used in combination with other therapies such as other chemotherapy, surgical treatment, radiation, thermotherapy, etc.

25 According to the fourth aspect of the present invention, a use of an LAK activity enhancer for obtaining a drug used for treating tumor and/or cancer is provided.

 In order to obtain a drug used for treating tumor and/or cancer, LAK activity enhancers of the present invention may be used in combination with other materials

having an immunopotentiating activity and antitumor activity and/or anticancer activity.

BRIEF DESCRIPTION OF THE DRAWINGS

5 FIG. 1 is a bar graph showing the LAK activity-enhancing effect of the extract of Lentinus edodes mycelium directly added to peripheral blood as in comparison with IL-2.

FIG. 2 is a bar graph showing the change of LAK
10 activity with the concentration of the extract of Lentinus edodes mycelium directly added to peripheral blood.

FIG. 3 is a bar graph showing the LAK activity obtained by directly adding Krestin to peripheral blood.

FIG. 4 is a bar graph showing the LAK activity in
15 peripheral blood before and after oral administration of the extract of Lentinus edodes mycelium.

THE MOST PREFERRED EMBODIMENTS OF THE INVENTION

An extract of Lentinus edodes mycelium contained in
20 LAK activity enhancers of the present invention refers to an extract obtained by crushing and decomposing mycelia grown from Lentinus edodes cultured on a solid medium, or a solid medium itself containing Lentinus edodes mycelia in the presence of water and an enzyme.

25 An extract of Lentinus edodes mycelium used herein is preferably obtained by, but not specifically limited to, the following process.

Lentinus edodes spawn is inoculated on a solid medium

based on bagasse (sugar cane residue) and defatted rice bran to grow mycelia, and then the solid medium containing the grown mycelia is delignified to enable about 30% by weight or less to pass through a 12-mesh sieve.

- 5 To this delignified solid medium are added water and one or more enzymes selected from cellulase, protease or glucosidase while maintaining said solid medium at a temperature of around 30-55°C, and said solid medium is crushed and ground in the presence of said enzyme so that
- 10 at least 70% by weight of bagasse fiber is able to pass through a 12-mesh sieve. Then, the temperature is raised to 80-100°C to ensure inactivation of the enzyme and sterilization, and the resulting suspension is filtered to give an extract of *Lentinus edodes* mycelium.
- 15 The extract of *Lentinus edodes* mycelium as prepared above may be directly used in LAK activity enhancers or LAK activity-enhancing formulations of the present invention, but conveniently concentrated and freeze-dried into powder to be stored and used in various forms. The freeze-dried
- 20 product is a brown powder with hygroscopic characteristics and has a peculiar taste and odor.

 LAK activity enhancers containing the prepared extract of *Lentinus edodes* mycelium (or the extract of *Lentinus edodes* mycelium on its own) may be used in LAK

25 therapy in place of IL-2 to enhance the immune activity against tumor of the subject.

 LAK therapy generally comprises the steps of culturing in vitro lymphocytes from the subject with IL-2

to induce LAK cells from the lymphocytes, and reinfusing the LAK cells into the subject. At the step of reinfusing LAK cells into the subject, IL-2 may be (directly) concurrently administered.

- 5 LAK activity enhancers containing the extract of *Lentinus edodes* mycelium of the present invention may be cultured in vitro with lymphocytes from the subject (human, animals, etc.) to induce LAK cells or may be (directly) concurrently administered to the subject at the step of
- 10 reinfusing the cultured lymphocytes to the subject. They may be used in combination with other antitumor and/or anticancer agents.

- LAK activity enhancers containing the extract of *Lentinus edodes* mycelium of the present invention may be
- 15 directly added to peripheral blood collected from the subject, but preferably sterilized with acetone before being added.

- LAK activity enhancers containing the extract of *Lentinus edodes* mycelium of the present invention can be
- 20 directly added to peripheral blood collected from the subject preferably at a concentration expressed as the extract of *Lentinus edodes* mycelium of $1\text{ }\mu\text{g} - 1\text{ mg}/10^6$ lymphocytes, more preferably $10\text{ }\mu\text{g} - 100\text{ }\mu\text{g}/10^6$ lymphocytes, most preferably $10\text{ }\mu\text{g} - 50\text{ }\mu\text{g}/10^6$ lymphocytes. The dose
- 25 per ml of peripheral blood, which partially depends on the level of lymphocytes in peripheral blood, approximately represents $1\text{ }\mu\text{g} - 2.5\text{ mg}$, preferably $10\text{ }\mu\text{g} - 250\text{ }\mu\text{g}$, more preferably $10\text{ }\mu\text{g} - 125\text{ }\mu\text{g}$.

LAK activity-enhancing formulations comprising an LAK activity enhancer containing the extract of Lentinus edodes mycelium of the present invention can be directly administered to the subject to enhance the anti-tumor immune activity of the subject, thereby providing an effect comparable to the effect elicited by LAK therapy using IL-2.

Induction of LAK cells in the subject such as a human or animals by directly administering the LAK activity enhancers of the present invention may make it possible to reduce physical stress associated with the isolation of a great number of leukocytes and the high cost resulting from producing mass culture. In addition, the use of LAK activity-enhancers of the present invention can also reduce the risk of contamination during the isolation of blood from the subject and reinfusion can also be reduced.

LAK activity-enhancing formulations of the present invention are administered most preferably, but are not limited to, the oral route in view of the least stress imposed on the subject and the like. They may also be administered via intravenous, intracutaneous, subcutaneous or intramuscular injection, or percutaneous, nasal, enteral or other routes.

Suitable dosage forms of LAK activity-enhancing formulations of the present invention include, but are not limited to, tablets, capsules, powders, granules, solutions, syrups, injections, creams, ointments, patches, sprays, suppositories, etc.

Pharmaceutically acceptable carriers that can be

optionally mixed with LAK activity-enhancing formulations of the present invention include, but not limited to, excipients such as lactose, dextrose, starch, crystalline cellulose; binders such as starch, gelatin, methyl
5 cellulose, polyvinylpyrrolidone; disintegrators such as starch, calcium carboxymethylcellulose, carboxymethyl starch; lubricants such as talc, stearates; coating agents such as sucrose, talc, gelatin; as well as various brighteners, flavoring agents, colorants, corrigents,
10 solubilizers, stabilizers, suspending agents, absorbefaciants or the like known in the art depending on the purpose. For use as injections, various diluents commonly used in this field of art such as water or ethyl alcohol can be used.

15 The route of administration, dose, frequency or others of LAK activity-enhancing formulations of the present invention are determined taking into account the age, weight, condition and other factors of the subject.

The extract of *Lentinus edodes* mycelium contained in
20 LAK activity-enhancing formulations of the present invention is highly safe as has been traditionally ingested as a component of food. Therefore, the dose is not be strictly limited. For example, the extract of *Lentinus edodes* mycelium is normally administered preferably at a
25 dose of 100 mg - 10000 mg 2-3 times daily, more preferably 500 mg - 5000 mg three times daily, most preferably 1000 mg - 1500 mg three times daily. It may be administered in combination with other drugs such as antitumor and/or

anticancer agents such as immunopotentiators, antibiotics, analgesics, antiinflammatory agents, etc.

LAK activity-enhancing formulations of the present invention can also be provided in the form of a food.

5 Preferred forms of food include powders, granules, pastes, jellies, etc. Granules desirably are supplemented with sugars such as lactose to add a sweet taste. LAK activity-enhancing formulations of the present invention can also be provided in the form of a drink. These foods or drinks may
10 be supplemented with vitamins, minerals such as calcium, alcohols, deodorants such as polyphenols in addition to the extract of *Lentinus edodes* mycelium. These foods or drinks may include the categories of specific health foods, medical foods or the like.

15 LAK activity-enhancing formulations of the present invention can also be provided in the form of a feed or feed additive. LAK activity-enhancing formulations of the present invention can be used as a feed or feed additive for domestic animals to treat and/or prevent tumor
20 occurring in the domestic animals or to treat and/or prevent bacterial or viral infections in domestic animals.

As a result, the amount of therapeutic agents such as antibiotics currently used can be reduced, thereby reducing farming costs. Another advantage is that the period during
25 which shipment of animals is suspended due to the administration of antibiotics can be shortened.

LAK activity enhancement with LAK activity enhancers of the present invention can be confirmed by, but not

limited to, the procedure below according to the method of Takagi et al. (Clinical Immunology, 19:245-249, 1987).

LAK activity test

LAK activity can be determined by ^{51}Cr release assay, [^3H] uridine assay or the like. In terms of convenience and objectivity, the ^{51}Cr release assay is preferably used.

The ^{51}Cr release assay is one of methods for in vitro determining the cytotoxicity against target cells of LAK cells induced from lymphocytes treated with an LAK activity enhancer, and comprises the steps of:

(i) adding ^{51}Cr -labeled sodium chromate to the target cells to label the target cells,

(ii) reacting the labeled target cells with induced LAK cells, and

(iii) measuring the amount of ^{51}Cr released into the cell culture supernatant from the target cells bursted by the LAK cells.

Subcultured cells used as target cells in the ^{51}Cr release assay are preferably Daudi cells or Raji cells. Target cells cultured in a culture flask or the like are collected by centrifugation, and labeled with ^{51}Cr via incubation in a 5% CO_2 incubator at 37°C for 1-2 hours after 100-150 μCi ^{51}Cr labeled-sodium chromate has been added. Culture media for target cells are those suitable for the growth of the cells used, and include, for example, liquid media such as RPMI 1640 or F-12 appropriately supplemented with serum, antibiotics, etc.

Cultured target cells are washed with PBS three times, and then suspended in a culture medium at a density of 1×10^6 cells/ml to serve for assay.

To induce LAK cells, lymphocytes are isolated from
5 peripheral blood of the subject. Heparin is added to
peripheral blood from the subject, and monocytes at the
interface are separated by density gradient centrifugation
on Ficoll-Conray solution (s.g. = 1.077). The separated
monocytes are washed with PBS (pH 7.4, without Ca and Mg)
10 2-3 times and then suspended in a culture medium
(preferably, RPMI 1640 medium (Gibco) containing FBS
(inactivated fetal bovine serum) and/or antibiotics, if
desired) at a density of 1×10^6 cells /ml. This
suspension is transferred to a Petri dish which has been
15 precoated with autoserum (plasma) at 37°C for 15 minutes,
and incubated at 37°C for 1 hour. Unattached cells are
recovered as lymphocyte fractions.

The lymphocytes as prepared above are suspended in a
culture medium at a predetermined concentration, preferably
20 1×10^5 - 10^6 cells/ml.

The extract of Lentinus edodes mycelium is added to
the medium as used for preparing the lymphocyte-suspending
solution at a final concentration of about 1-100 µg/ml.

Predetermined amounts of the lymphocyte-suspending
25 solution and extract-containing medium are added to wells
of a culture plate. In parallel, predetermined amounts of
the lymphocyte-floating solution and the medium not
containing the extract of Lentinus edodes mycelium are

added to wells as an extract-free control. Then, the culture plate is statically incubated in a CO₂ incubator at room temperature for 3 days.

When a 96-well U-bottom microtest plate is used, for example, 100 µl of the lymphocyte-floating solution is preferably added to each well at a density of 1×10^4 to 1×10^5 cells/well. The number of cells per well can be appropriately determined by those skilled in the art in accordance with the well volume, the activity of LAK cells, the sensitivity of target cells to LAK cells, etc.

Thus, lymphocytes derived from the subject are cultured in the presence of the extract of *Lentinus edodes* mycelium of the present invention at various concentrations (including zero) to induce effector cells. As used herein, the term effector cells refers to cells subjected to said culture treatment for 3 days and includes both lymphocytes cocultured with an LAK activity enhancer (lymphocytes treated with an LAK activity enhancer) and lymphocytes cultured in the medium alone without LAK activity enhancer (lymphocytes treated under the extract-free condition).

Cultured lymphocytes (effector cells) are recovered from the well and resuspended in fresh RPMI 1640 medium containing 10% FBS (1×10^6 cells/ml).

In the assay for determining cytotoxicity, each well containing said target cells is further supplied with a predetermined amount of the floating solution containing said various effector cells (lymphocytes treated with LAK active enhancer or those under the extract-free condition)

at a density of 1×10^5 - 1×10^6 cells/ml for experimental dissociation, 1N HCl in an amount equal to the effector-floating solution for maximum dissociation, or the medium alone in an amount equal to the effector-floating solution for natural dissociation. Then, cells are collected at the bottom of the well of the culture plate using a plate centrifuge or the like, and then incubated in a 5% CO₂ incubator at 37°C for a predetermined period (for example, 2-5 hours).

After completion of the incubation, the radioactivity of ⁵¹Cr released in the culture supernatant in the culture supernatant in each well is measured using a scintillation counter or the like.

The term "maximum dissociation" refers to the total amount of ⁵¹Cr released when all the target cells are destroyed. Maximum dissociation can be determined by, for example, lysing target cells via incubation in a culture medium containing 1N HCl under predetermined conditions in the absence of effector cells and measuring the amount of ⁵¹Cr released in the culture supernatant.

The term "experimental dissociation" refers to the amount of ⁵¹Cr released by cytotoxicity of effector cells against target cells and natural death of target cells when effector cells are added to target cells. Experimental dissociation can be determined by, for example, incubating target cells under the same conditions as those for maximum dissociation with the exception of adding effector cells in place of 1N HCl to the culture medium and measuring the

amount of ^{51}Cr released in the culture supernatant.

The term "natural dissociation" refers to the amount of ^{51}Cr released by natural death of target cells. Natural dissociation can be determined by, for example, incubating target cells under the same conditions as for maximum dissociation with exception of containing neither 1N HCl nor effector cells and measuring the amount of ^{51}Cr released in the culture supernatant.

The activation level of LAK cells is evaluated on the basis of the LAK activity calculated by equation (1) below.

$$\text{LAK activity\%} = \frac{\text{Experimental dissociation(cpm)} - \text{Natural dissociation(cpm)}}{\text{Maximum dissociation(cpm)} - \text{Natural dissociation(cpm)}} \times 100$$

Equation (1)

The disclosure of JPA Nos. 353927/98 and 353928/98 on which the present application is based to claim the priority are incorporated herein as a whole as reference.

The following examples further illustrate the present invention but should not be taken as limiting the scope of the invention thereto. It will be appreciated by those skilled in the art that various changes and modifications can be made without departing from the spirit of the present invention.

EXAMPLES

Example 1: Preparation of an extract of *Lentinus edodes* mycelium

A solid medium consisting of 90 parts by weight of

bagasse and 10 parts by weight of rice bran was soaked with an appropriate amount of pure water, and then inoculated with *Lentinus edodes* spawn and allowed to stand in an incubator at controlled temperature and humidity to grow mycelia. After mycelia spread over the solid medium, the bagasse base was delignified to enable 24% by weight or less to pass through a 12-mesh sieve. To 1.0 kg of this delignified medium were added 3.5 L of pure water and 2.0 g of purified cellulase while maintaining the solid medium at 40°C to prepare a medium-containing mixture.

Then, the medium-containing mixture was circulated by a variable speed gear pump, during which the solid medium was crushed and ground at the gears for about 200 minutes so that about 80% by weight of bagasse fiber is able to pass through a 12-mesh sieve. The medium-containing mixture was crushed and ground while the temperature of said mixture was gradually increased. Then, the medium-containing mixture was further heated to 90°C to ensure inactivation of the enzyme and sterilization and allowed to stand at 90°C for 30 minutes. The resulting medium-containing mixture was filtered through a 60-mesh filter cloth to give an extract of *Lentinus edodes* mycelium solution, which was concentrated and then converted into a freeze-dried powder.

The extract of *Lentinus edodes* mycelium as prepared above contained 25.3% (w/w) carbohydrates determined by the phenol-sulfuric acid method, 19.7% (w/w) proteins determined by the Lowry method and 2.6% (w/w) polyphenols

determined by the Folin-Denis method using gallic acid as standard. The extract of Lentinus edodes mycelium further contains 8% crude fat, 22% crude ash and about 20% soluble nitrogen-free materials other than carbohydrates.

- 5 The extract of Lentinus edodes mycelium had a sugar composition (%) as follows: Xyl 15.2, Ara 8.2, Man 8.4, Gul 39.4, Gal 5.4, GlcN 12.0, GluUA 11.3.

10 Example 2: Comparison of LAK activities of the extract of Lentinus edodes mycelium and IL-2

 The extract of Lentinus edodes mycelium obtained in Example 1 was directly administered to peripheral blood collected from the subject to compare the LAK activity-enhancing effect with that of IL-2.

- 15 Heparin was added to peripheral blood from subject A, and monocytes at the interface were separated by density gradient centrifugation on Ficoll-Conray solution (s.g. = 1.077). The separated monocytes were washed with PBS (pH 7.4, without Ca and Mg) twice and then suspended in RPMI 20 1640 medium (Gibco) containing 10% FBS (inactivated fetal bovine serum) at a density of 1×10^6 cells /ml. This cell suspension was transferred to a Petri dish which had been precoated with autoserum (plasma) at 37°C for 5 minutes followed by incubation at 37°C for 1 hour, and then 25 unattached cells were recovered as lymphocyte fractions.

 The lymphocytes as prepared above were suspended in RPMI 1640 medium (Gibco) containing 10% FBS (inactivated serum) at 1×10^6 cells /ml.

03050710.012002

LAK activity enhancer-containing solutions were prepared by adding IL-2 at a final concentration of 25 U/ml or the extract of *Lentinus edodes* mycelium obtained in Example 1 at final concentrations of 1, 10, 100 or 1000
5 µg/ml to RPMI 1640 medium (Gibco) containing 10% FBS (inactivated serum).

To each well of a 96-well U-bottom microtest plate were added 100 µl of the lymphocyte-suspending solution and 100 µl of IL-2 or the LAK activity enhancer-containing
10 solution containing the extract of *Lentinus edodes* mycelium at various concentrations (a total of 200 µl). As an extract-free control, 100 µl of the lymphocyte-suspending solution and 100 µl of the extract-free medium (RPMI 1640 medium (Gibco) containing 10% FBS (inactivated serum)) were
15 added. Then, the culture plate was statically incubated in a CO₂ incubator at room temperature for 3 days.

Cultured lymphocytes (effector cells) were recovered from the well and resuspended in fresh RPMI 1640 medium containing 10% FBS at a concentration of 1×10^6 cells/ml.

20 Target cells (Daudi cells from Dainippon Pharmaceutical) subcultured in RPMI 1640 medium containing 10% FBS were recovered by centrifugation, and incubated with 100-150 µCi/ 10^6 cells of ⁵¹Cr labeled-sodium chromate (New England Nuclear) in a 5% CO₂ incubator at 37°C for 1
25 hour. Cultured cells labeled with ⁵¹Cr were washed with PBS three times, and then suspended in RPMI 1640 medium containing 10% FBS at a concentration of 1×10^6 cells/ml.

A 50 µl aliquot (5×10^4 cells/well) of labeled

target cells was added to each of wells of the microtest plate for maximum dissociation, natural dissociation and experimental dissociation. In addition, 100 μ l of 1N HCl was added to each well for maximum dissociation, 100 μ l of
5 RPMI 1640 medium containing 10% FBS was added to each well for natural dissociation, and 100 μ l of effector cells treated with 10 μ g/ml of the extract of *Lentinus edodes* mycelium of the present invention at concentrations of 1, 10, 100 and 1000 μ g/ml or 25 U/ml of IL-2 or the medium
10 alone were added to each well for experimental dissociation. Here, each well for experimental dissociation, maximum dissociation and natural dissociation was not located around the outer periphery of the plate.

Then, the microtest plate was centrifuged at 800 rpm
15 for 5 minutes on a plate centrifuge to collect cells at the bottom of the well, and then incubated in a 5% CO₂ incubator at 37°C for 3.5 hours.

The culture supernatant in each well was collected by SOKEN-PET Σ -96 from the incubated plate, and the
20 radioactivity was measured in a γ -scintillation counter.

LAK activity was calculated by equation (1) above.

The results are shown in Table 1 and Fig. 1.

Table 1

Test No.	Inducer	LAK activity
1	None	11%
2	IL-2 (final concentration: 25 U/ml)	55%
3	Extract of Lentinus edodes mycelium (final concentration: 1 µg/ml)	14%
4	Extract of Lentinus edodes mycelium (final concentration: 10 µg/ml)	20%
5	Extract of Lentinus edodes mycelium (final concentration: 100 µg/ml)	14%

In order to examine the toxicity of the extract of Lentinus edodes mycelium to lymphocytes, lymphocytes was treated with the extract of Lentinus edodes mycelium as prepared in Example 1 at a final concentration of 1 mg/ml by the same procedure as described above and observed under a microscope to show that almost 100% of lymphocytes survived (data not shown).

Example 3: Optimal concentration of the extract of Lentinus edodes mycelium for LAK activity

The optimal concentration of the extract of Lentinus edodes mycelium to be directly added to peripheral blood collected from the subject was examined. LAK activity was calculated in the same manner as described in Example 2 with exception that the extract of Lentinus edodes mycelium as prepared in Example 1 was added at final concentrations of 1, 5, 10 and 100 µg/ml to peripheral blood collected

from subjects B and C.

The results are shown in Table 2 and Fig. 2.

As shown in Table 2 and Fig. 2, the extract of
Lentinus edodes mycelium of the present invention shows the
5 highest LAK activity when it is added at a final
concentration of 10 $\mu\text{g/ml}$ or 50 $\mu\text{g/ml}$ (i.e. 10 μg or 50 μg
per 10^6 lymphocytes). The dose to peripheral blood also
depends on the proportion of lymphocytes in peripheral
blood of the subject, but approximately corresponds to
10 about 10-25 μg or 50-125 μg of the extract of Lentinus
edodes mycelium per 1 ml of peripheral blood.

Table 2

Test No.	Inducer	LAK activity	
		Subject	
		B	C
1	None	13%	27%
2	Extract of Lentinus edodes mycelium (final concentration: 1 $\mu\text{g/ml}$)	18%	28%
3	Extract of Lentinus edodes mycelium (final concentration: 5 $\mu\text{g/ml}$)	19%	29%
4	Extract of Lentinus edodes mycelium (final concentration: 10 $\mu\text{g/ml}$)	21%	34%
5	Extract of Lentinus edodes mycelium (final concentration: 50 $\mu\text{g/ml}$)	19%	36%
6	Extract of Lentinus edodes mycelium (final concentration: 100 $\mu\text{g/ml}$)	15%	30%

Comparative example 1: LAK activity of Krestin

LAK activity-enhancing effect of an immunopotentiator Krestin was examined to compare it with the effect of the extract of *Lentinus edodes* mycelium of the present

5 invention. LAK activity-enhancing effect of Krestin was evaluated in the same manner as described in Example 2 with exception that Krestin (from Kureha Chemical Industry) was added to peripheral blood collected from subject D at a final concentration of 1 mg/ml.

10 The final concentration of 1 mg/ml is the concentration said to be the most effective when Krestin is used to induce LAK cells.

The results are shown in Table 3 and Fig. 3.

As shown in Table 3 and Fig. 3, the LAK-enhancing
15 activity of Krestin was lower than that of a control not containing LAK activity enhancer even at the optimal concentration of 1 mg/ml.

Table 3

Test No.	Inducer	LAK activity
1	None	14%
2	Krestin (final concentration: 1 mg/ml)	13%

20

Example 4: Determination of LAK activity before and after administration of a formulation containing the extract of *Lentinus edodes* mycelium

An LAK activity-enhancing formulation containing the
25 extract of *Lentinus edodes* mycelium prepared in the same

manner as described in Example 1 was administered to subjects B, C, E and F to evaluate the LAK activity-enhancing effect.

This LAK activity-enhancing formulation comprises 0.6 g of the extract of *Lentinus edodes* mycelium and 2.4 g of lactose per 3 g of the formulation, and was administered at a dose of 6 g (1.2 g as the extract of *Lentinus edodes* mycelium) three times daily or a total of 3.6 g/day expressed as the extract of *Lentinus edodes* mycelium for a week. Peripheral blood was collected from each subject before administration and about 2 hours after the final administration, and LAK activity was determined in the same manner as described in Example 2.

The results are shown in Table 4 and Fig. 4.

As shown in Table 4 and Fig. 4, the extract of *Lentinus edodes* mycelium of the present invention remarkably increased LAK activity in all of subjects B, C, E and F.

20

Table 4

Subject	<u>LAK activity</u>			
	B	C	E	F
Before administration of LEM	13%	27%	14%	18%
After administration of LEM	40%	43%	18%	28%

In order to examine side effects caused by the administration of the extract of *Lentinus edodes* mycelium, biochemical blood tests were performed on subjects B, C and

E. The results are shown in Table 5. Throughout the period for administration of the extract of *Lentinus edodes* mycelium, subjects B, C, E and F did not show any side effects, such as generalized fatigue, chills, fever or
5 digestive system symptoms.

As also shown in Table 5, any side effects were not observed by administration of the extract of *Lentinus edodes* mycelium of the present invention.

Table 5

	Subject B		Subject C		Subject E	
	Administration for 1 week					
	Before	After	Before	After	Before	After
WBC (x10 ³ /μl)	5.45	5.94	5.85	5.08	8.01	7.47
RBC (x10 ⁶ /μl)	4.56	4.61	4.99	4.85	4.89	4.65
Hb (g/dl)	14.2	14.2	14.4	13.8	15.5	14.8
Ht (%)	42.2	42.8	43.6	42.1	44.8	42.9
MCV (fl)	92.5	92.8	87.4	86.8	91.6	92.3
MCH (pg)	31.1	30.8	28.9	28.5	31.7	31.8
MCHC (g/dl)	33.6	33.2	33.0	32.8	34.6	34.5
PLT (x10 ³ /μl)	19.1	18.5	23.9	19.3	29.2	29.7
RDW-SD (fl)	43.7	44.6	43.5	43.6	43.7	44.8
RDW-CV (%)	12.8	13.0	13.4	13.6	13.0	13.3
PDW (fl)	14.0	13.4	11.6	12.6	13.3	13.2
MPV (fl)	11.1	10.9	10.0	10.2	10.6	10.4
P-LCR (%)	34.1	32.5	24.9	27.6	30.1	28.8
Stab	0.5	0.5	1.0	0.0	0.5	0.0
Seg	54.0	48.5	56.0	43.5	46.5	38.5
Lym	37.5	40.5	38.0	43.5	28.5	35.0
Mono	4.5	7.5	3.0	9.5	11.5	8.5
Eo	2.5	3.0	1.5	1.0	12.0	15.5
Baso	1.0	0.0	0.5	0.5	0.5	0.5
TP (g/dl)	6.6	6.9	7.7	7.5	8.0	7.7
ALB (g/dl)	3.9	4.1	4.4	4.3	4.8	4.7
A/G	1.44	1.46	1.33	1.34	1.50	1.57
TTT (SHU)	3.9	5.0	8.0	8.1	5.3	3.0
ZTT (Kunk)	8.6	8.4	12.6	12.4	6.3	4.9
U-N (mg/dl)	10.4	10.3	10.2	11.2	11.8	11.6
U-A (mg/dl)	5.6	5.9	5.4	5.2	6.1	5.9
Creatinine (mg/dl)	0.67	0.71	0.84	0.82	0.67	0.72
Overall BIL (mg/dl)	0.6	0.7	0.5	0.6	0.5	0.5
Direct BIL (mg/dl)	0.1	0.1	0.0	0.0	0.1	0.1
Indirect BIL (mg/dl)	0.5	0.6	0.5	0.6	0.4	0.4
GLU (mg/dl)	124	104	94	111	115	114
AST (GOT) (IU/l)	15	13	18	16	33	33
ALT (GPT) (IU/l)	29	30	20	20	45	57
LD (LDH) (IU/l)	142	139	171	152	380	343
CK (CPK) (IU/l)	121	131	227	177	317	224
ALP (IU/l)	131	136	156	148	174	162
γ-GT (IU/l)	12	14	23	23	225	240
LAP (IU/l)	45	40	80	73	134	120
CHE (IU/l)	2358	2593	2779	2749	3191	3299
AMY (IU/l)	78	87	107	93	59	54
T-CHO (mg/dl)	208	232	206	200	255	243
HDLCHO (mg/dl)	50	50	44	40	47	47
T-G (mg/dl)	135	210	270	303	451	359
Na (mEq/l)	141	142	140	141	139	141
K (mEq/l)	3.7	3.9	4.1	3.8	4.0	4.2
Cl (mEq/l)	106	107	105	107	103	104
Ca (mEq/l)	4.4	4.4	4.6	4.4	4.9	4.7
I-P (mg/dl)	3.3	3.1	3.0	2.7	3.5	3.2
H	0	0	0	0	1	0
L	0	0	0	1	1	0
I	0	1	1	1	1	1
RA test	(-)	(-)	(-)	(-)	(-)	(-)
CRP (mg/dl)	0.03	0.06	0.02	0.01	0.07	0.12
ANA	<40	<40	<40	<40	<40	<40
ANA pattern	(-)	(-)	(-)	(-)	(-)	(-)
CH50 (CH50/ml)	33.0	33.0	45.0	42.0	53.0	52.0

INDUSTRIAL APPLICABILITY

LAK enhancers containing the extract of Lentinus edodes mycelium of the present invention showed similar effects to those of IL-2 with no direct toxicity to lymphocytes when they were directly administered to peripheral blood. They are available at far lower cost as compared with IL-2. Therefore, they can be substituted for expensive IL-2 which also has the disadvantage of causing strong side effects.

Formulations containing the extract of Lentinus edodes mycelium of the present invention could remarkably increase LAK activity with no side effects even by direct administration. Therefore, formulations containing the extract of Lentinus edodes mycelium of the present invention are useful as an immunopotentiator capable of increasing LAK activity with a benign side-effect profile by direct administration such as oral administration without using LAK therapy.

CLAIMS

1. An LAK activity enhancer containing an extract of *Lentinus edodes* mycelium.
2. The LAK activity enhancer of claim 1 for enhancing LAK activity by acting on lymphocytes derived from peripheral blood.
3. The LAK activity enhancer of claim 2 containing an extract of *Lentinus edodes* mycelium at a concentration of 1 μg or more per 10^6 lymphocytes.
4. An LAK activity-enhancing formulation containing the LAK activity enhancer of claim 1.
5. A pharmaceutical or veterinary LAK activity-enhancing formulation comprising the LAK activity enhancer of claim 1 and a pharmaceutically acceptable carrier.
6. The LAK activity-enhancing formulation of claim 4 or 5 for oral administration.
7. The LAK activity-enhancing formulation of claim 4 in the form of a food, drink or feed.
8. The LAK activity-enhancing formulation of claim 4 or 5 for injection or percutaneous administration.
9. The LAK activity-enhancing formulation of any one of claims 4 to 8, wherein said formulation is used for treating tumor and/or cancer.
10. A method for treating tumor and/or cancer comprising administering an effective amount of the LAK activity-enhancing formulation of any one of claims 4 to 8.
11. A use of the LAK activity enhancer of claim 1 for the preparation of a drug used for treating tumor and/or

cancer.

2022081409562

ABSTRACT

It is an object of the present invention to provide immunopotentiators and immunotherapeutic formulations available at low cost and having a benign side effect profile. According to the present invention, an LAK activity enhancer containing an extract of *Lentinus edodes* mycelium and an LAK activity-enhancing formulation containing said enhancer are provided.

FIG. 1




FIG. 2




FIG. 3





FIG. 4



BIRCH, STEWART, KOLASCH & BIRCH, LLP

PLEASE NOTE:
YOU MUST
COMPLETE THE
FOLLOWING:

COMBINED DECLARATION AND POWER OF ATTORNEY

ATTORNEY DOCKET NO

0230-0161P

FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:
LAK ACTIVITY ENHANCERS DERIVED FROM EXTRACT OF LENTINUS EDODES MYCELIUM AND
LAK ACTIVITY-ENHANCING FORMULATIONS CONTAINING THE EXTRACT

Insert Title:

Fill in Appropriate
Information -
For Use Without
Specification
Attached:

the specification of which is attached hereto. If not attached hereto,

the specification was filed on _____ as
United States Application Number _____; and /or

the specification was filed on November 26, 1999 as PCT
International Application Number PCT/JP99/06616; and was
amended under PCT Article 19 on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Insert Priority
Information:
(if appropriate)

→ Prior Foreign Application(s)

<u>353927/1998</u>	<u>Japan</u>	<u>11/27/98</u>
(Number)	(Country)	(Month/Day/Year Filed)
<u>353928/1998</u>	<u>Japan</u>	<u>11/27/98</u>
(Number)	(Country)	(Month/Day/Year Filed)
_____	(Country)	(Month/Day/Year Filed)
(Number)	(Country)	(Month/Day/Year Filed)
_____	(Country)	(Month/Day/Year Filed)
(Number)	(Country)	(Month/Day/Year Filed)

Priority Claimed

<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No

Insert Provisional
Application(s):
(if any)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)
_____	_____

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More Than 12 Months (6 Months for Designs) Prior To The Filing Date of This Application:

Insert Requested
Information:
(if appropriate)

Country	Application No	Date of Filing (Month/Day/Year)
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Insert Prior U.S.
Application(s):
(if any)

(Application Number)	(Filing Date)	(Status - patented, pending, abandoned)
_____	_____	_____
(Application Number)	(Filing Date)	(Status - patented, pending, abandoned)
_____	_____	_____

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Terrell C. Birch (Reg. No. 19,382)
Joseph A. Kolasch (Reg. No. 22,463)
Bernard L. Sweeney (Reg. No. 24,448)
Charles Gorenstein (Reg. No. 29,271)
Leonard R. Svensson (Reg. No. 30,330)
Andrew D. Meikle (Reg. No. 32,868)
Joe McKinney Muncy (Reg. No. 32,334)
C. Joseph Faraci (Reg. No. 32,950)

Raymond C. Stewart (Reg. No. 21,066)
James M. Slattery (Reg. No. 23,380)
Michael K. Mutter (Reg. No. 29,689)
Gerald M. Murphy, Jr. (Reg. No. 28,977)
Terry L. Clark (Reg. No. 32,644)
Marc S. Weiner (Reg. No. 32,181)
Andrew F. Reish (Reg. No. 33,443)
Donald J. Daley (Reg. No. 34,313)

Send Correspondence to:

BIRCH, STEWART, KOLASCH & BIRCH, LLP

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:
YOU MUST
COMPLETE THE
FOLLOWING:

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First or Sole
Inventor:
Insert Name of Inventor
Insert Date This
Document is Signed

Insert Residence
Insert Citizenship

Insert Post Office
Address

Full Name of Second
Inventor, if any:

see above

Full Name of Third
Inventor, if any

see above

Full Name of Fourth
Inventor, if any

see above

Full Name of Fifth
Inventor, if any

see above

GIVEN NAME <u>Kenji</u>	FAMILY NAME <u>ASANO</u>	INVENTOR'S SIGNATURE <u>Kenji Asano</u>	DATE* May 30, 2001
Residence (City, State & Country) <u>Osaka, Japan</u>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <u>c/o Kobayashi Pharmaceutical Co., Ltd. of 3-13-35, Mitsuya Minami, Yodogawa-ku, Osaka-shi, Osaka 532-0035 Japan</u>			
GIVEN NAME <u>Yukiko</u>	FAMILY NAME <u>MATSUDA</u>	INVENTOR'S SIGNATURE <u>Yukiko Matsuda</u>	DATE* May 30, 2001
Residence (City, State & Country) <u>Osaka, Japan</u>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <u>c/o Kobayashi Pharmaceutical Co., Ltd. of 3-13-35, Mitsuya Minami, Yodogawa-ku, Osaka-shi, Osaka 532-0035 Japan</u>			
GIVEN NAME <u>Yutaka</u>	FAMILY NAME <u>TAJIMA</u>	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country) <u>Saga, Japan</u>		CITIZENSHIP <u>Japanese</u>	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country) <u>10-511, Yaemizo 3-chome, Saga-shi, Saga 849-0935 Japan</u>			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			

* DATE OF SIGNATURE